




Original Research

Carbapenem-resistant *Klebsiella pneumoniae*: Risk Factors for Isolation Among Hospitalized Patients

Ketevan Kobaidze, MD¹ , Jesse Jacob, MD², Traci Leong, PhD³, W Dana Flanders, MD³

¹ Emory Division of Hospital Medicine, Emory Department of Medicine, Emory University School of Medicine, ² Emory Division of Infectious Diseases, Emory Department of Medicine, Emory University School of Medicine, ³ Department of Biostatistics and Bioinformatics. Rollins School of Public Health, Emory University

Keywords: Carbapenem resistant *Klebsiella pneumoniae*, Multi drug resistance, antibiotic stewardship, carbapenem-resistant Enterobacterales, CRE <https://doi.org/10.56305/001c.57568>

Vol. 2, Issue 1, 2023

Background: Carbapenem-resistant *Klebsiella pneumoniae* (CRKP) is an important healthcare-associated pathogen. This study aimed to identify factors associated with CRKP isolation among hospitalized patients, describe molecular epidemiology, and mortality associated with CRKP isolation. **Methods:** We performed a case-control study at two university-affiliated teaching hospitals. We included 150 patients (30 cases and 120 controls). Each patient with CRKP, a case-patient, was matched with four controls by admission facility, admission date, age, and sex. Controls, patients without CRKP, were randomly selected from a computerized list of inpatients whose admission date was the same as that of the case, within 48 hours of the date of the initial positive culture. We calculated the risk of in-hospital death as the number of deaths divided by the number of cases and evaluated the risk of mortality associated with the site of positive culture. Molecular epidemiology investigation using comparison of restricted DNA patterns of CRKP by pulsed-field gel electrophoresis (PFGE) was conducted. **Results:** A greater proportion of cases than controls had undergone an invasive procedure, including use of a central vein catheter, or mechanical nutrition by tube feeding. Pre-admission treatment within two months with the following antibiotic classes was associated with CRKP isolation: carbapenems, fluoroquinolones, anti-pseudomonal penicillins, and cephalosporins. The molecular analysis indicated that over 90% of isolates shared similar PFGE patterns. CRKP isolation was associated with significantly higher in-hospital mortality in comparison to controls. Positive cultures from sites other than urine were associated with substantially higher mortality than was a positive urine culture (RR= 4.0). **Conclusions:** The use of multiple broad-spectrum antibiotics, multiple comorbid conditions and poor performance status are important risk factors for developing CRKP in a hospitalized population.

INTRODUCTION

For the past decade, national surveillance studies have documented a growing rate of antimicrobial resistance in hospitals in the United States.¹ Infection caused by multidrug-resistant Gram-negative bacteria present a challenge to clinicians since it is associated with increased mortality, morbidity, length of hospital stays, and financial cost.² The mode of acquisition and spread of resistant Gram-negative bacilli is complex. These organisms are highly efficient at acquiring multidrug-resistant coding genes under selective antibiotic pressure and are capable of rapid dissemination.^{3,4}

The global rise in antimicrobial resistance coupled with the limited new anti-microbial agents increasingly drove treatment choices toward carbapenems. Carbapenems, the most enzyme-stable class of β -lactam antibiotics, have been considered the last resort antimicrobial therapy against many antibiotic-resistant Gram-negative organisms.

Carbapenem-resistant *Klebsiella pneumoniae* (CRKP) has emerged as a serious healthcare-associated pathogen in the United States and worldwide.³⁻⁶ CRKP threatens hospitalized patients and lengthens recovery since very limited therapeutic options are available to treat this infection. Large city-wide outbreaks have been reported in northeastern states, where it became endemic.

The first CRKP isolate in the Emory Healthcare system was detected in September 2006. The patient was from New York City, an endemic area for CRKP, seeking evaluation for a wound infection following liver transplantation. Because there were frequent transfers within the Emory Healthcare system (two 500-bed acute care hospitals, 1 long-term acute care facility (LTAC), 2 rehabilitation units, and 1 geriatric hospital), there was a potential for spread.

We conducted a case-control study to examine factors associated with isolation of CRKP among the hospitalized population in the two acute care hospitals, Emory University Hospital Midtown (EUHM) and Emory University Hospital (EUH).

METHODS

All adult patients aged ≥ 18 years who had been treated in the hospital from September 2006 through December 2008 and had a clinical culture positive for CRKP were enrolled. A case-patient was identified once the initial CRKP culture was reported and included only once, even if CRKP was isolated again.

Each case-patient was matched with four control-patients by admission facility, age ± 5 years, sex, and date of culture, as a time-at-risk for hospitalized patients.

Controls, patients without CRKP, were randomly selected from a computerized list of inpatients who matched the case age (± 5 years), sex, and facility and whose admission date were within 48 hours of the date of the initial, positive culture. Patients, who spent less than 48 hours in the hospital, and obstetric service patients were not included as controls. We chose the control group from the population at risk which allowed us to estimate the risk associated with specific exposures, such as prior use of antimicrobial agents.⁷ It also allowed us to compare the mortality associated with CRKP isolation to that without CRKP. A health-care exposure was defined as hospitalization of more than 48h, previous hospitalization within six months, previous LTAC or nursing home placement, or hemodialysis.

From the medical record, we abstracted information including basic demographics, admission date, outcome, services, and admission facilities, urinary catheter use, central line placement, intubation, and mechanical ventilation before CRKP isolation, antibiotic use within two months before hospitalization as it was documented in the chart, or electronic records and in the hospital prior CRKP isolation; exposure to interventional procedures, tube feeding, and length of stay in the hospital. The performance status was measured by the Karnofsky scale per physical and occupational services evaluation at the time of CRKP isolation.

The Charlson comorbidity score was used to assess and compare patients' co-morbidities, and APACHE-II scores were used for those who were treated in the ICU.⁸

Laboratory confirmation of CRKP was restricted to detection of *bla*_{-KPC} gene by PCR.⁹ For PCR, the bacterial DNA was extracted using the EZ-1 BioRobot (Qiagen, Germantown, MD). Amplification and detection of a 399 base pair segment of the *bla*_{-KPC} gene were performed on the 7500 Real-Time PCR system (Applied Biosystems, Foster City, CA). All isolates were found to contain the *bla*_{-KPC} gene by PCR. Molecular subtyping by PFGE was performed to detect genetic relatedness among isolates.

Descriptive analyses were conducted, including means, medians, and percentiles. The comparison of clinical characteristics between cases and controls was made using Chi-Square (or Fisher's exact) tests. Relative risks (RR), and exact confidence intervals (CI) for the association between risk factors and CRKP were estimated using exact conditional logistic regression (Proc Exact, in SAS, version v9.2). Conditional logistic regression was used to account for our matched design. Exact distributions were used due to the small numbers for some combinations of exposure and dis-

ease. In separate analyses of the cases, we calculated the risk of in-hospital death as the number of deaths divided by the number of cases. To assess the association of mortality with the site of culture, we calculated the risk ratio: the proportion who died among those for whom CRKP was isolated from a site other than urine divided by the proportion who died among those for whom CRKP was isolated from the urine.

Molecular subtyping by pulsed-field gel electrophoresis (PFGE) was performed for the comparison of restricted DNA patterns of all CRKP isolates.¹⁰ For better molecular characterization additional 21 isolates from all sites (total 51 isolates) were included in molecular subtyping including our case - patients 30 isolates.

This study was approved by the Institutional Review Board of the Emory University School of Medicine.

RESULTS

STUDY POPULATION

The first case was identified in September of 2006, 13 cases were detected in 2007, and 16 cases in 2008 across these two hospitals (total of 30 with 120 matched controls). No seasonal variations were apparent in the identification of CRKP.

The clinical characteristics of patients with CRKP enrolled in this study (summarized in [Table 1](#)), show the median age was 60 years. Cultures were obtained from blood, urine, various wounds, and respiratory tract (upper or lower tract). Urine was the most common site of organism isolation, followed by lung and wound. Of 30 cases eleven patients died (36.7%) whereas 4 patients in the control group (3.3%) died. Nine of the 11 deceased cases had CRKP isolated from specimens other than urine or multiple sites. Cases with a positive culture from a site other than urine had substantially higher mortality than patients whose positive cultures were only from urine (RR= 4.0). Four patients had received solid organ transplants before CRKP isolation, and two of those died in the hospital.

All but one patient was confirmed to have healthcare-acquired CRKP. The source of CRKP acquisition in one patient was not identified and was considered to be non-healthcare related

RISK FACTORS ASSOCIATED WITH ISOLATION OF CRKP

Factors assessed for association with isolation of CRKP in the hospital are summarized in [Table 2](#). A greater proportion of cases than controls had undergone an invasive procedure, including the use of a central vein catheter ($p=0.007$, OR, 3.4, 95% CI, 1.4-8.7), or mechanical ventilation ($p=0.002$, OR, 3.6, 95% CI, 1.6-8.1). Nutrition by tube feeding ($p=0.001$, OR, 4.2, 95% CI, 1.8 -10) and admission from a nursing home or LTAC were also associated with increased risk ($p<0.001$, OR=6.2, 95% CI, 2.5-15.1).

Most patients with CRKP had been treated with two or more antibiotics within 2 months prior to the current hospitalization ($p<0.005$, OR, 8.5, 95% CI, 3.3 -21.7). Pre-admission treatment two months before hospitalization with

Table 1. Case-patients characteristics

Total number of patients (%)	30 (100%)
LOS, median	28
Admission service	
Medicine	22 (73)
Surgery (or other)	8 (27)
Age, years	
Mean	60
Median	62 (range from 27 to 90)
Sex	
Female	25 (52)
Male	23 (48)
Site of isolation	
Urine	15 (50)
Blood	6 (13)
Sputum /BAL	4 (21)
Wound /other	5 (17)
Survived	19
Died in the hospital	11
Survivals disposition	
Home	10 (51)
NH/LTAC	6(32)
Other	3(17)
Nosocomial acquisition	29 (97)
Non-nosocomial acquisition	1 (3)

the following antibiotic classes was associated with CRKP acquisition: carbapenems ($p=0.001$, OR, 24.4, 95% CI, 2.73-217.96), fluoroquinolones ($p<0.001$, OR, 6.17, 95% CI, 2.4 – 15.83), anti-pseudomonal penicillin ($p<0.02$, OR, 6.03, 95% CI, 1.98 -18.32), and cephalosporins ($p = 0.001$, OR, 5.36, 95% CI, 2.07 -13.87).

Post-admission treatment with anti-pseudomonal penicillins ($p<0.001$, OR, 6.5, 95% CI, 2.7-15.4), carbapenems ($p=0.016$, OR, 12.4, 95% CI, 1.59 -100), and a fluoroquinolone ($p<0.03$, OR, 2.6, 95% CI, 1.1-5.7) were significantly associated with CRKP isolation. Interestingly cephalosporin antibiotic use in the hospital was not a significant risk factor (Table 2).

On average, cases stayed in the hospital 32.5 days compared to 10 days for controls, and the average time from admission to first CRKP isolation was 12.3 days. A Charlson score of seven or higher ($p=0.047$, OR, 1.12, 95% CI, 1.0 – 1.26), and low scores of performance status by Karnofsky scale ($p<0.001$, OR, 0.93, 95% CI, 0.91- 0.96) were significantly associated with the isolation of CRKP.

MOLECULAR SUBTYPING

The PFGE pattern analysis (Fig 1) showed that over 90% of the isolates shared similar PFGE patterns (>85% band similarity). In addition, the prevailing pattern was found to be very similar to PFGE patterns of isolates that the Centers for Disease Control (CDC) determined to be multilocus sequential type ST258, and which appears to be the dominant CRKP strain seen throughout the United States.¹¹

DISCUSSION

Since its isolation in 1996 in North Carolina, CRKP has been detected in multiple states and across the world including China, Europe, Central and South America, and Israel.¹¹⁻¹⁷ It is a challenging infection with attributable mortality as high as 40%, and potential to rapidly spread. CRKP is now endemic in the Northeastern United States, particularly in the New York area, but periodic outbreaks have been reported throughout the country.¹⁸⁻²² Some authors speculate that the global spread could be greater than reported since the detection of CRKP by clinical microbiology laboratories remains difficult.

Previous studies have identified the inter-hospital spread of CRKP and the results of our study also suggest that CRKP was introduced in our system by an index case in September 2006, and subsequently spread through these hospitals.¹⁸⁻²⁰ Molecular sub-typing by PFGE confirmed >85% genetic relationship among the isolates. Our results are consistent with previous reports that CRKP dissemination occurs in part by the spread of isolates from patients previously hospitalized in an endemic area.^{11,23} Two of our patients traveled from New York City and sought medical attention in our facilities.

Even though the number of incidents remained low we observed a significant increase in incidence during the study period. Detection of CRKP infection likely represents only part of the problem since the vast majority of transmissions likely occur through colonized asymptomatic patients. Infection spread was well documented in this study, showing how a strain of CRKP can quickly disseminate throughout a hospital.

Most prior studies of risk factors for CRKP acquisition have focused on antimicrobial exposure, underlying conditions, ICU exposure, and procedures.^{5,6,24-28} The present study confirmed the use of multiple (two or more) antibiotics, including cephalosporins, fluoroquinolones, anti-pseudomonal penicillins, and carbapenems, before hospitalization was associated with increased risk isolation of CRKP. Previous research by Gasink and colleagues had identified that fluoroquinolone use can be a risk factor for the acquisition of CRKP which is consistent with the results of our study.²⁹

The *bla*-*KPC* gene, responsible for resistance, resides on a transmissible plasmid and its relocation from other *Enterobacteriaceae* species has been confirmed by phage-typing.³⁰ Mathers, et al. have shown several mechanisms by which the CRKP resistance gene can spread including plasmid transfer and clonal spread.²⁴ We found the occurrence of CRKP- producing *Klebsiella oxytoca* and *Citrobacter freundii* isolates in addition to CRKP in two of our cases. We were not able to perform phage typing to confirm the resistance gene interspecies translocation.

Hussein and colleagues detected that exposure to at least one antibiotic drug before isolating CRKP was a risk factor.²⁰ In contrast, our study identified that pre-hospital use of two or more antibiotics within two months significantly increased the risk of isolation of CRKP, whereas the

Table 2. Risk factors associated with isolation of CRKP among hospitalized patients

	Case	Control	P-Value	Odds Ratio	Confidence Interval
Antibiotics (pre- hospitalization)			<.001		
0	11 (36.7%)	100 (81.3%)			
1	4 (13.3%)	7 (5.7%)	.493	1.639	(.39,6.76)
2+	15 (50%)	16 (13%)	<.001	8.547	(3.3,21.74)
Antipseudomonal PCN	8 (26.7%)	7 (5.7%)	.002	6.026	(1.98,18.32)
Cephalosporines	11 (36.7%)	12 (9.8%)	.001	5.355	(2.07,13.87)
Vancomycin	2 (6.7%)	9 (7.3%)	1.0	.905	(.19,4.42)
Quinolones	12 (40%)	12 (9.8%)	<.0001	6.167	(2.40,15.83)
Carbapenem	5 (16.7%)	1 (.85)	.001	24.4	(2.73,217.96)
Antibiotics (during hospitalization)**			.005		
0	1 (3.3%)	18 (14.6%)			
1	5 (16.7%)	37 (30.1%)	0.99	0	(1.29, 10.64)
2+	24 (80%)	48 (39%)	.015	3.7	(1.78, 29.31)
Carbapenem	15 (50%)	4 (3.3%)	.016	12.5	(1.59,100)
Cephalosporines	4 (13.3%)	39 (31.7%)	.068	.331	(.11,1.02)
Aminoglycosides	2 (6.7%)	6 (4.9%)	.655	1.393	(.27,7.27)
Anti-pseudomonal PCN	20 (66.7%)	29 (23.6%)	<.001	6.483	(2.73,15.41)
Vancomycin	5 (16.7%)	16 (13.0%)	.565	1.34	(.45, 3.99)
Quinolones	16 (53.3%)	38 (30.9%)	.032	2.556	(1.13,5.76)
Intubated/Vent	16 (53.3%)	30 (24.4%)	0.002	3.55	(1.55, 8.13)
Total Length of Stay	32.5 (38.6)	10 (10.2)	0.001	1.029	(1.01,1.05)
Charlson Score	7.4 (4.3)	6 (3.3)	0.047	1.123	(1.00,1.26)
Urinary Catheter	24 (80%)	85 (69.1%)	0.270	1.79	(0.68, 4.74)
APACHE II for ICU	32	10	0.2	1.027	(.986, 1.069.)
Central Vein Catheter	23 (76.7%)	60 (48.8%)	0.007	3.44	(1.38, 8.62)
Tube Feeding	14 (46.7%)	21 (17.1%)	0.001	4.26	(1.80, 10)
Pre-hospital Site					
NH + LTAC + Hospital	22 (73.3%)	38 (30.9%)	<.001	6.151	(2.51, 15.06)
Home	8 (26.7%)	85 (69.1%)			
KPS score	39.3	60.4	<.001	0.934	(0.91, 0.96)

Legends: LOS-length of stay, NH – Nursing home; LTAC – long-term-acute-care facility
PCN=penicillin, KPS – Karnofsky performance status

Charlson score, APACHE II, and KPS scores indicate mean value

** For cases, represents antibiotic use prior to CRKP isolation

use of one antibiotic before hospitalization didn't reach a significant difference.

The results of the present study add to a substantial body of data demonstrating the role of previous antibiotic use in the development CRKP mechanism. Our findings show that pre-hospital use of carbapenems was associated with a higher risk of acquiring CRKP. We also found that not only carbapenems but fluoroquinolones and anti-pseudomonal antibiotic use were associated with significant risk while patients were treated in the hospital.

Interestingly, in-hospital cephalosporins use was not associated with a CRKP isolation risk, while within two months of pre-hospital use, this antibiotic was associated with a significantly increased risk of CRKP isolation. Patients in our study had longer hospitalizations and underwent more interventional procedures including central line placement and intubation with mechanical ventilation,

than controls. Case patients were treated in the ICU more often than controls (data not shown), APACHE II scores for those who were admitted to the ICU detected higher numbers of cases than controls but the difference was not statistically significant (Table 2).

Previous studies suggested that underlying multiple medical conditions were associated with a higher risk of acquisition of CRKP.²⁷⁻²⁹ Our results support the conclusion that multiple comorbidities, here measured by the Charlson score, are a risk factor.

Hospitalized immunocompromised patients, including solid organ transplant recipients, are at increased risk to suffer from infection with multidrug-resistant pathogens.²⁴ Consistent with this finding is that four cases patients enrolled in our study were solid organ recipients, two of whom died in the hospital.




Figure 1. Dendrogram of similarity of PFGE patterns.

Isolates above #2009728543 has 85% similar band patterns, 5 isolates pattern has distinct patterns. ID# 2009728542 -the index case isolate from 2006

Although the increased risk for urinary tract infection and use of indwelling bladder catheter is well documented in CRKP, the study detected that most of our patients with this catheter had no greater risk for CRKP than without it.^{15,31} This finding was explained by the frequent use of indwelling catheters among all hospitalized patients.

CRKP outbreaks have also been reported in LTAC and nursing homes.^{31–33} The present study identified admission from LTAC or nursing home as a risk factor associated with isolation of CRKP. This observation further supports the previous finding that CRKP is truly a healthcare-associated infection. Only one case-patient was admitted from their home, without identifiable exposure to healthcare. Although the number is small, it suggests that CRKP can be acquired in the community

Our study had several limitations. No surveillance screening was performed before this study to identify the CRKP carrier state initially. Due to the small number of cases, we decided to include all patients with CRKP, thus all patients documented to have infection or colonization were enrolled. Also due to the small sample size, we accommodated only for the matching factor, but not other covariates thus only univariate analysis was performed. The study was conducted in tertiary teaching hospitals close to LTAC, our

finding cannot be generalized to all CRKP cases for other hospitals.

These findings have several important clinical implications. The use of multiple broad-spectrum antibiotics remains an important risk factor for developing CRKP. Moreover, hospitalized patients with multiple comorbidities and poor performance status are susceptible to CRKP, thus this risk group should be spared from unnecessary antibiotic exposure, and judicious use of antibiotics should be used by healthcare providers.

Consideration should be given when performing multiple invasive procedures since invasive procedures are associated with a higher risk for CRKP isolation.

CRKP infection poses a significant threat to the already vulnerable hospitalized patients. The most common carbapenemases in the United States are *K. pneumoniae* carbapenemases (KPCs).

The treatment of antimicrobial-resistant infections will continue to challenge clinicians. As newer antibiotics against resistant pathogens are incorporated into clinical practice, we are learning more about their effectiveness and propensity to resistance. Treatment guidelines from the Infectious Disease Society of America are available.⁵

Efforts aimed at understanding and updating the epidemiology of CRKP among hospitalized patients should continue.

.....

CORRESPONDING AUTHOR

Ketevan Kobaidze, MD, Ph.D.
Emory University School of Medicine
550 Peachtree Street, Atlanta, GA, 30308
Phone: 1-404 840 4395
Fax: 1- 404 686 4837
Email: kkobaid@emory.edu

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

FUNDING STATEMENT

Supported by Emory University School of Medicine Research Grant

AUTHOR CONTRIBUTIONS

All authors have reviewed the final manuscript prior to submission. All the authors have contributed significantly to the manuscript, per the International Committee of Medical Journal Editors criteria of authorship.

- Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; AND
- Drafting the work or revising it critically for important intellectual content; AND
- Final approval of the version to be published; AND
- Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

ACKNOWLEDGMENTS

None.

Submitted: September 06, 2022 EST, Accepted: December 09, 2022 EST



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc/4.0> and legal code at <https://creativecommons.org/licenses/by-nc/4.0/legalcode> for more information.

REFERENCES

1. Kallen AJ, Hidron AI, Patel J, Srinivasan A. Multidrug resistance among gram-negative pathogens that caused healthcare-associated infections reported to the national healthcare safety network, 2006–2008. *Infect Control Hosp Epidemiol*. 2010;31(5):528–531. doi:10.1086/652152
2. Peleg AY, Hooper DC. Hospital-acquired infections due to gram-negative bacteria. *N Engl J Med*. 2010;362(19):1804–1813. doi:10.1056/nejmra0904124
3. Nordmann P, Cuzon G, Naas T. The real threat of *Klebsiella pneumoniae* carbapenemase-producing bacteria. *Lancet Infect Dis*. 2009;9(4):228–236. doi:10.1016/s1473-3099(09)70054-4
4. Calfee D, Jenkins SG. Use of active surveillance cultures to detect asymptomatic colonization with carbapenem-resistant *Klebsiella pneumoniae* in intensive care unit patients. *Infect Control Hosp Epidemiol*. 2008;29(10):966–968. doi:10.1086/590661
5. Falagas ME, Rafailidis PI, Kofteridis D, et al. Risk factors of carbapenem-resistant *Klebsiella pneumoniae* infections: a matched case control study. *J Antimicrob Chemother*. 2007;60(5):1124–1130. doi:10.1093/jac/dkm356
6. Samra Z, Ofir O, Lishtzinsky Y, Madar-Shapiro L, Bishara J. Outbreak of carbapenem-resistant *Klebsiella pneumoniae* producing KPC-3 in a tertiary medical centre in Israel. *Int J Antimicrob Agents*. 2007;30(6):525–529. doi:10.1016/j.ijantimicag.2007.07.024
7. Harris AD, Samore MH, Lipsitch M, Kaye KS, Perencevich E, Carmeli Y. Control-group selection importance in studies of antimicrobial resistance: example applied to *Pseudomonas aeruginosa*, *Enterococci*, and *Escherichia coli*. *Clin Infect Dis*. 2002;34(12):1558–1563. doi:10.1086/340533
8. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chron Dis*. 1987;40(5):373–383. doi:10.1016/0021-9681(87)90171-8
9. Cole JM, Schuetz AN, Hill CE, Nolte FS. Development and evaluation of a real-time PCR assay for detection of *Klebsiella pneumoniae* carbapenemase genes. *J Clin Microbiol*. 2009;47(2):322–326. doi:10.1128/jcm.01550-08
10. Tenover FC, Arbeit RD, Goering RV, et al. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J Clin Microbiol*. 1995;33(9):2233–2239. doi:10.1128/jcm.33.9.2233-2239.1995
11. Kitchel B, Sundin DR, Patel JB. Regional dissemination of KPC-producing *Klebsiella pneumoniae*. *Antimicrob Agents Chemother*. 2009;53(10):4511–4513. doi:10.1128/aac.00784-09
12. Wei ZQ, Du XX, Yu YS, Shen P, Chen YG, Li LJ. Plasmid-mediated KPC-2 in a *Klebsiella pneumoniae* isolate from China. *Antimicrob Agents Chemother*. 2007;51(2):763–765. doi:10.1128/aac.01053-06
13. Centers for Disease Control and Prevention. Vital signs: carbapenem-resistant enterobacteriaceae. *MMWR Morb Mortal Rep*. 2013;62(early release):1–6.
14. Naas T, Nordmann P, Vedel G, Poyart C. Plasmid-mediated carbapenem-hydrolyzing beta-lactamase KPC in a *Klebsiella pneumoniae* isolate from France. *Antimicrob Agents Chemother*. 2005;49(10):4423–4424. doi:10.1128/aac.49.10.4423-4424.2005
15. Villegas MV, Lolans K, Correa A, et al. First detection of the plasmid-mediated class A carbapenemase KPC-2 in clinical isolates of *Klebsiella pneumoniae* from South America. *Antimicrob Agents Chemother*. 2006;50(8):2880–2882. doi:10.1128/aac.00186-06
16. Rossi F. The challenges of antimicrobial resistance in Brazil. *Clin Infect Dis*. 2011;52(9):1138–1143. doi:10.1093/cid/cir120
17. Navon-Venezia S, Chmelnitsky I, Leavitt A, Schwaber MJ, Schwartz D, Carmeli Y. Plasmid-mediated imipenem-hydrolyzing enzyme KPC-2 among multiple carbapenem-resistant *Escherichia coli* clones in Israel. *Antimicrob Agents Chemother*. 2006;50(9):3098–3101. doi:10.1128/aac.00438-06
18. Patel G, Huprikar S, Factor SH, Jenkins SG, Calfee DP. Outcomes of carbapenem-resistant *Klebsiella pneumoniae* infection and the impact of antimicrobial and adjunctive therapies. *Infect Control Hosp Epidemiol*. 2008;29(12):1099–1106. doi:10.1086/592412
19. Bratu S, Landam D, Haag R, et al. Rapid spread of carbapenem-resistant *Klebsiella pneumoniae* in New York City: a new threat to our antibiotic armamentarium. *Arch Intern Med*. 2005;165:1430–1435.

20. Hussein K, Sprecher H, Mashiach T, Oren I, Kassis I, Finkelstein R. Carbapenem resistance among *Klebsiella pneumoniae* isolates: risk factors, molecular characteristics, and susceptibility patterns. *Infect Control Hosp Epidemiol*. 2009;30(7):666-671. doi:10.1086/598244
21. Gupta N, Limbago BM, Patel JB, Kallen AJ. Carbapenem-resistant Enterobacteriaceae: epidemiology and prevention. *Clin Infect Dis*. 2011;53(1):60-67. doi:10.1093/cid/cir202
22. Howard-Anderson JR, Earley M, Komarow L, et al. Poor outcomes in both infection and colonization with carbapenem-resistant Enterobacteriales. *Infect Control Hosp Epidemiol*. Published online February 2, 2022:1-7. doi:10.1017/ice.2022.4
23. Adler A, Shklyar M, Schwaber MJ, et al. Introduction of OXA-48-producing Enterobacteriaceae to Israeli hospitals by medical tourism. *J Antimicrob Chemother*. 2011;66(12):2763-2766. doi:10.1093/jac/dkr382
24. Mathers AJ, Cox HL, Bonatti H, et al. Fatal cross infection by carbapenem-resistant *Klebsiella* in two liver transplant recipients. *Transpl Infect Dis*. 2009;11(3):257-265. doi:10.1111/j.1399-3062.2009.00374.x
25. Marchaim D, Navon-Venezia S, Schwaber MJ, Carmeli Y. Imipenem-resistant Enterobacter species isolation: the emergence of KPC-none carbapenemase, molecular characterization, epidemiology, and outcomes. *Antimicrob Agents Chemother*. 2008;52(4):1413-1418. doi:10.1128/aac.01103-07
26. Giakoupi P, Maltezou H, Polemis M, Pappa O, Saroglou G, Vatopoulos A. KPC-2-producing *Klebsiella pneumoniae* infections in Greek hospitals are mainly due to a hyperepidemic clone. *Euro Surveill*. 2009;14(21):pii=19218.
27. Schwaber MJ, Klarfeld-Lidji S, Navon-Venezia S, Schwartz D, Leavitt A, Carmeli Y. Predictors of carbapenem-resistant *Klebsiella pneumoniae* acquisition among hospitalized adults and effect of the acquisition on mortality. *Antimicrob Agents Chemother*. 2008;52(3):1028-1033. doi:10.1128/aac.01020-07
28. Kwak YG, Choi SH, Choo EJ, et al. Risk factors for the acquisition of carbapenem-resistant *Klebsiella pneumoniae* among hospitalized patients. *Microb Drug Resist*. 2005;11(2):165-169. doi:10.1089/mdr.2005.11.165
29. Gasink LB, Zaoutis TE, Bilker WB, Lautenbach E. The categorization of prior antibiotic use: impact on the identification of risk factors for drug resistance in case control studies. *Am J Infect Control*. 2007;35(10):638-642. doi:10.1016/j.ajic.2007.01.011
30. Rasheed JK, Biddle JW, Anderson KF, et al. Detection of the *Klebsiella pneumoniae* carbapenemase Type 2 carbapenem-hydrolyzing enzymes in clinical isolates of *Citrobacter freundii* and *K. oxytoca* carrying a common plasmid. *J Clin Microbiol*. 2008;46(6):2066-2069. doi:10.1128/jcm.02038-07
31. Carbapenem-resistant *Klebsiella pneumoniae* associated with a long-term care facility - West Virginia, 2009-2011. *MMWR*. 2011;60:1418-1420.
32. Chitnis AS, Caruthers PS, Rao AK, et al. Outbreak of carbapenem-resistant Enterobacteriaceae at a long-term acute care hospital. *Infect Control Hosp Epidemiol*. 2012;33(10):984-992. doi:10.1086/667738
33. Mills J, Chapin K, Andrea S, Furtado G, Mermel L. Community and nursing home residents with carbapenemase-producing *Klebsiella pneumoniae* infection. *Infect Control Hosp Epidemiol*. 2011;32(6):629-631. doi:10.1086/660202